

# LINEAR IDENTITIES IN GRAPH ALGEBRAS

AGATA PILITOWSKA

ABSTRACT. We find the basis of all linear identities which are true in the variety of entropic graph algebras. We apply it to describe the lattice of all subvarieties of power entropic graph algebras.

## 1. INTRODUCTION

In 1979 Caroline Shallon introduced in her dissertation [9] algebras associated with graphs. Let  $G = (V, E)$  be a (undirected) graph with a set  $V$  of vertices and a set  $E \subseteq V \times V$  of edges. Its *graph algebra*  $A(G) = (V \cup \{0\}, \cdot)$  is a groupoid with the multiplication defined as follows:

$$x \cdot y := \begin{cases} x, & \text{if } (x, y) \in E, \\ 0, & \text{otherwise.} \end{cases}$$

C. Shallon proved that many finite graph algebras are nonfinitely based. Moreover it was shown in [4] that a graph algebra  $A(G)$  is finitely based if and only if it is entropic.

In this paper we find all linear identities which are true in the variety of entropic graph algebras. Some new linear identity (3.1) true in this variety is crucial for the final result. Linear identities play an important rôle in the theory of power algebras.

The *power* (complex or global) *algebra*  $\mathbf{CmA} = (\mathcal{P}(A), F)$  of an algebra  $(A, F)$  is the family  $\mathcal{P}(A)$  of all non-empty subsets of  $A$  with complex operations given by

$$f(A_1, \dots, A_n) := \{f(a_1, \dots, a_n) \mid a_i \in A_i\},$$

where  $\emptyset \neq A_1, \dots, A_n \subseteq A$  and  $f \in F$  is an  $n$ -ary operation.

Power algebras were studied by several authors, for instance by G. Grätzer and H. Lakser [5], S. Whitney [6], A. Shafaat [8], C. Brink [3], I. Bošnjak and R. Madarász [2].

Complex operations may preserve some of properties of  $(A, F)$ , but not all identities true in  $(A, F)$  must be satisfied in  $\mathbf{CmA}$ . For example, the power algebra of a group is not again a group [5]. For an arbitrary variety  $\mathcal{V}$ , G. Grätzer and H. Lakser determined the identities satisfied by the variety generated by  $\mathbf{CmA}$ , for  $\mathbf{A} \in \mathcal{V}$ . They applied their result to describe all subvarieties of power algebras of lattices and groups [5]. They showed that there is exactly one non-trivial variety of

---

2000 *Mathematics Subject Classification.* Primary 08B15, 03C05, 03C13, 08C10; Secondary 17D99, 08A05, 08A40, 08A62, 08A99.

*Key words and phrases.* graph algebra, linear identity, entropic algebra, equational basis, lattice of subvarieties, power algebra of subsets.

While working on this paper, the author was partially supported by the INTAS grant #03-51-4110 and Statutory Grant of Warsaw University of Technology # 504G11200013000.

power algebras of lattices and there are exactly three non-trivial varieties of power algebras of groups.

In this paper we show that the lattice of all subvarieties of the variety generated by power algebras of entropic graph algebras has eight elements.

In Section 2 we recall main known results concerning graph algebras. Section 3 is devoted to linear identities satisfied in entropic graph algebras. In the last Section 4 we present some theorems concerning identities satisfied by power algebras of sets and describe all subvarieties of power entropic graph algebras.

We say that an algebra  $(A, F)$  is *idempotent* if each element  $a \in A$  forms a one-element subalgebra of  $(A, F)$ . An algebra  $(A, F)$  is *entropic*, if each operation  $f \in F$  as a mapping from a direct power of the algebra into the algebra is actually a homomorphism.


We call a term  $t$  *linear*, if every variable occurs in  $t$  at most once. An identity  $t \approx u$  is called *linear*, if both terms  $t$  and  $u$  are linear. An identity  $t \approx u$  is called *regular*, if  $t$  and  $u$  contain the same variables. The notation  $t(x_1, \dots, x_n)$  means that the term  $t$  contains no other variables than  $x_1, \dots, x_n$  (but not necessarily all of them).

In the case of groupoid terms we will use non-brackets notation, as follows:

$$xx_1x_2 \dots x_n := \begin{cases} (\dots((xx_1)x_2)\dots)x_n, & n \geq 1 \\ x, & n = 0 \end{cases}$$

## 2. GRAPH ALGEBRAS

For each natural number  $n \in \mathbb{N}$ , let  $P_n$  denote  $n$ -vertex graphs in the form of a path without loops, and  $L_n$  denote  $n$ -vertex graphs in the form of a path with loops, as in the diagrams:



For graphs  $G$  and  $H$ ,  $G + H$  will denote the disjoint union of  $G$  and  $H$ , with no edges between  $G$  and  $H$ .

It is not difficult to see that the direct product of two graph algebras is not necessarily a graph algebra. So the class of all graph algebras does not form a variety. Let us denote by  $\mathcal{V}_{A(G)}$  the variety generated by a graph algebra  $A(G)$ .

As was shown in [1], the variety of all entropic graph algebras is generated by the algebra  $A(P_2 + L_2)$ . The following identities

$$(2.1) \quad xy \approx xyy$$

$$(2.2) \quad x(yz) \approx xy(yz)$$

$$(2.3) \quad xyz \approx xzy$$

$$(2.4) \quad xy \approx x(yx)$$

$$(2.5) \quad x(yzu) \approx x(yz)(yu)$$

$$(2.6) \quad x(y(zu)) \approx x(yz)(uz)$$

$$(2.7) \quad x(yz)(uv) \approx x(yv)(uz)$$

$$(2.8) \quad x(xy) \approx x(yy)$$

$$(2.9) \quad xx(yz) \approx x(yy)(zz)$$

form a basis for  $\mathcal{V}_{A(P_2+L_2)}$ . It is easy to see that the entropic law

$$(E) \quad xy(zw) \approx xz(yw).$$

follows by the above identities:

$$xy(zw) \stackrel{(2.4)}{\approx} x(yx)(zw) \stackrel{(2.7)}{\approx} x(yw)(zx) \stackrel{(2.3)}{\approx} x(zx)(yw) \stackrel{(2.4)}{\approx} xz(yw).$$

The lattice of all subvarieties of the variety of entropic graph algebras was discussed in [1] and it is given in Figure 1.




FIGURE 1

$\mathcal{T}$  denotes the trivial variety,  $\mathcal{SL}$  is the variety of semilattices,  $\mathcal{LZ}$  is the variety of left zero bands (groupoids determined by identity  $xy \approx x$ ),  $\mathcal{LN}$  is the variety of left normal bands - idempotent semigroups satisfying the additional left-normal law (2.3),  $\mathcal{U}_1$  is the variety defined by two identities: (2.1) and

$$(2.10) \quad xx \approx xy.$$

and  $\mathcal{U}_2$  is the variety defined by all identities (2.1)-(2.9) and additional one

$$(2.11) \quad xx \approx x(yy).$$

The remaining subvarieties of entropic graph algebras have the following basis:

$$\begin{aligned}
\mathcal{V}_{A(P_1)} & : xy \approx uz; \\
\mathcal{V}_{A(P_1+L_1)} & : x(yz) \approx (xy)z, xy \approx yx, xy \approx x(yy); \\
\mathcal{V}_{A(P_1+L_2)} & : (2.1) - (2.9), x(yz) \approx (xy)z; \\
\mathcal{V}_{A(P_2)} & : (2.1) - (2.9), xx \approx yy, x(yz) \approx z(yx); \\
\mathcal{V}_{A(P_2+L_1)} & : (2.1) - (2.9), x(yz) \approx z(yx); \\
\mathcal{V}_{A(P_3)} & : (2.1) - (2.9), xx \approx yy; \\
\mathcal{V}_{A(P_3+L_1)} & : (2.1) - (2.9), x(yy) \approx y(xx).
\end{aligned}$$

As it was proved in [1] using the regular identities (2.1)-(2.9), every term in the variety  $\mathcal{V}_{A(P_2+L_2)}$  may be expressed in one of the standard forms:

- i)  $x$ ,
  - ii)  $x_1(x_1x_1)(x_2x_2) \dots (x_nx_n)$ , for  $n \geq 1$ ,
  - iii)  $x_1(y_1x_1)(y_2x_2) \dots (y_nx_n) \approx (x_1y_1)(y_2x_2) \dots (y_nx_n)$ ,
- for  $n \geq 1$ , where  $\{x_1, \dots, x_n\} \cap \{y_1, \dots, y_n\} = \emptyset$ . The variables  $x_1, \dots, x_n$  will be referred to as bottom and  $y_1, \dots, y_n$  will be referred to as top variables.

**Theorem 2.1.** [1] *An identity  $p \approx q$  is derivable from identities (2.1)-(2.9) if and only if both sides of the identity  $p \approx q$  have the same standard form, the same leftmost variables and (in the case of type (iii)) the same top and bottom variables.*

### 3. LINEAR IDENTITIES IN ENTROPIC GRAPH ALGEBRAS

In this section we will find all linear identities which are true in the variety  $\mathcal{V}_{A(P_2+L_2)}$ . First we prove that all entropic graph algebras satisfy some linear identity which plays an important rôle in this paper.

**Theorem 3.1.** *The following linear identity*

$$(3.1) \quad x(y(zt)) \approx xy(tz)$$

is satisfied in the variety  $\mathcal{V}_{A(P_2+L_2)}$ .

*Proof.*

$$\begin{aligned}
x(y(zt)) & \stackrel{(2.6)}{\approx} x(yz)(tz) \stackrel{(2.2)}{\approx} xy(yz)(tz) \\
& \stackrel{(E)}{\approx} xyt(yzz) \stackrel{(2.1)}{\approx} xyt(yz) \stackrel{(E)}{\approx} xy(yz)(tz) \stackrel{(2.1)}{\approx} xy(tz).
\end{aligned}$$

□

Let  $\Sigma$  be the following set of identities:

$$\Sigma : \begin{cases} (E) & xy(zt) \approx xz(yt), \\ (2.3) & xyz \approx xzy, \\ (3.1) & x(y(zt)) \approx xy(tz). \end{cases}$$

These identities are all linear and regular and they hold in any variety of entropic graph algebras. Moreover, the linear identity (2.7) also follows from  $\Sigma$ :

$$x(yz)(tw) \stackrel{(E)}{\approx} xt(yzw) \stackrel{(2.3)}{\approx} xt(ywz) \stackrel{(E)}{\approx} x(yw)(tz).$$

We will show that any linear identity true in  $\mathcal{V}_{A(P_2+L_2)}$  is a consequence of  $\Sigma$ . Before proving this fact we present a sequence of technical lemmas.

Let  $n \geq 1$  and  $\pi$  be any permutation of the set  $\{1, 2, \dots, n\}$ . By using identities from  $\Sigma$  repeatedly we obtain:

$$(3.2) \quad (xy_1y_2 \dots y_n)(zt_1t_2 \dots t_n) \approx xz(y_1t_1) \dots (y_nt_n), \text{ (by (E))}$$

$$(3.3) \quad xy_1y_2 \dots y_n \approx xy_{\pi(1)}y_{\pi(2)} \dots y_{\pi(n)}, \text{ (by (2.3))}$$

$$(3.4) \quad x(y(z_1t_1)(z_2t_2) \dots (z_nt_n)) \approx xy(t_1z_1)(t_2z_2) \dots (t_nz_n). \text{ (by (3.1))}$$

Moreover, as it was proved in [1], the following identity follows by entropicity and identities (2.3) and (2.7):

$$(3.5) \quad \begin{aligned} &xy_1 \dots y_k(y_{k+1}z_1) \dots (y_{k+n}z_n) \approx \\ &xy_{\pi(1)} \dots y_{\pi(k)}(y_{\pi(k+1)}z_{\delta(1)}) \dots (y_{\pi(k+n)}z_{\delta(n)}), \end{aligned}$$

where  $\pi$  is a permutation on the set  $\{1, 2, \dots, k+n\}$  and  $\delta$  is a permutation on the set  $\{1, 2, \dots, n\}$ .

**Lemma 3.2.** *The following identities*

$$(3.6) \quad xy_1 \dots y_n \approx x(y_1x)(y_2x) \dots (y_nx),$$

$$(3.7) \quad x(yz_1z_2 \dots z_n) \approx x(yx)(yz_1)(yz_2) \dots (yz_n)$$

hold in the variety  $\mathcal{V}_{A(P_2+L_2)}$  for any  $n \geq 1$ .

*Proof.* The proof of (3.6) goes by induction. For  $n = 1$ , the identity (3.6) follows by (2.4). Now let us assume that (3.6) is true for some  $n > 1$ . Then using (2.3) and (2.4) we obtain

$$\begin{aligned} xy_1 \dots y_{n+1} &\approx x(y_1x)(y_2x) \dots (y_nx)y_{n+1} \stackrel{(2.3)}{\approx} xy_{n+1}(y_1x)(y_2x) \dots (y_nx) \\ &\stackrel{(2.4)}{\approx} x(y_{n+1}x)(y_1x)(y_2x) \dots (y_nx) \stackrel{(2.3)}{\approx} x(y_1x)(y_2x) \dots (y_{n+1}x). \end{aligned}$$

The identity (3.7) follows directly by (3.6) and (3.4):

$$\begin{aligned} &x(yz_1z_2 \dots z_n) \stackrel{(3.6)}{\approx} x(y(z_1y)(z_2y) \dots (z_ny)) \\ &\stackrel{(3.4)}{\approx} xy(yz_1) \dots (yz_n) \stackrel{(2.4)}{\approx} x(yx)(yz_1) \dots (yz_n). \end{aligned}$$

□

**Lemma 3.3.** *Let  $n, k \geq 0$ . The following identities follow from the set  $\Sigma$ :*

$$(3.8) \quad x(y(t_1t_2z_1 \dots z_n)) \approx xyz_1 \dots z_n(t_2t_1),$$

$$(3.9) \quad (x(y_1y_2 \dots y_n))(t_1t_2z_1 \dots z_k) \approx (x(y_1y_2 \dots y_nz_1 \dots z_k))(t_1t_2),$$

$$(3.10) \quad (x(ty_1 \dots y_nw))z \approx (x(ty_1 \dots y_n))(zw),$$

$$(3.11) \quad (x(z_1y_1 \dots y_n))(z_2w) \approx (x(z_2y_1 \dots y_n))(z_1w),$$

$$(3.12) \quad \begin{aligned} &(yy_1y_2 \dots y_n)(zz_1z_2 \dots z_k) \approx \\ &\begin{cases} (y(zz_1 \dots z_{k-n}))(y_1z_{k-n+1}) \dots (y_nz_k), & \text{if } k > n \geq 0 \\ yy_1 \dots y_{n-k}z(y_{n-k+1}z_1) \dots (y_nz_k), & \text{if } n \geq k \geq 0 \end{cases} \end{aligned}$$

$$(3.13) \quad \begin{aligned} &(x(z_1y_1 \dots y_k))(z_2y_{k+1}) \dots (z_{n+1}y_{k+n}) \approx \\ &(x(z_{\pi(1)}y_{\delta(1)} \dots y_{\delta(k)}))(z_{\pi(2)}y_{\delta(k+1)}) \dots (z_{\pi(n+1)}y_{\delta(k+n)}), \end{aligned}$$

for any permutation  $\delta$  on the set  $\{1, 2, \dots, k+n\}$  and any permutation  $\pi$  on the set  $\{1, 2, \dots, n+1\}$ .

*Proof.* 1. For  $n = 0$ , the identity (3.8) follows by (3.1). If  $n > 0$ , then

$$x(y(t_1 t_2 z_1 \dots z_{n-1} z_n)) \stackrel{(3.1)}{\approx} (xy)(z_n(t_1 t_2 z_1 \dots z_{n-1})).$$

By induction on  $n$  and (3.3) we obtain

$$(xy)(z_n(t_1 t_2 z_1 \dots z_{n-1})) \approx xyz_n z_1 \dots z_{n-1}(t_2 t_1) \stackrel{(3.3)}{\approx} xy z_1 \dots z_n(t_2 t_1).$$

2. For  $n = 0$  and  $k = 0$ , the identity (3.9) is obvious. If  $n > 0$  and  $k > 0$ , then

$$\begin{aligned} (x(y_1 y_2 \dots y_n))(t_1 t_2 z_1 \dots z_{k-1} z_k) &\stackrel{(3.1)}{\approx} x(y_1 y_2 \dots y_n(z_k(t_1 t_2 z_1 \dots z_{k-1}))) \\ &\stackrel{(3.3)}{\approx} x(y_1(z_k(t_1 t_2 z_1 \dots z_{k-1}))y_2 \dots y_n). \end{aligned}$$

Using (3.8) we conclude that  $y_1(z_k(t_1 t_2 z_1 \dots z_{k-1})) \approx y_1 z_k z_1 \dots z_{k-1}(t_2 t_1)$ . Thus

$$\begin{aligned} x(y_1(z_k(t_1 t_2 z_1 \dots z_{k-1}))y_2 \dots y_n) &\approx x(y_1 z_k z_1 \dots z_{k-1}(t_2 t_1)y_2 \dots y_n) \\ &\stackrel{(3.3)}{\approx} x(y_1 y_2 \dots y_n z_1 \dots z_k(t_2 t_1)) \stackrel{(3.1)}{\approx} (x(y_1 y_2 \dots y_n z_1 \dots z_k))(t_1 t_2). \end{aligned}$$

3. For  $n = 0$ , the identity (3.10) follows by (2.3) and entropicity. If  $n > 0$ , then

$$(x(ty_1 \dots y_n w))z \stackrel{(2.3)}{\approx} (xz)(ty_1 \dots y_n w) \stackrel{(E)}{\approx} (x(ty_1 \dots y_n))(zw).$$

4. For  $n = 0$ , the identity (3.11) follows by entropicity. If  $n > 0$ , then

$$(x(z_1 y_1 \dots y_n))(z_2 w) \stackrel{(2.3)}{\approx} (x(z_2 w))(z_1 y_1 \dots y_n) \stackrel{(E)}{\approx} (x(z_1 y_1 \dots y_{n-1}))(z_2 w y_n).$$

By induction on  $n$ , (2.3) and (2.7) we obtain

$$\begin{aligned} (x(z_1 y_1 \dots y_{n-1}))(z_2 w y_n) &\approx (x(z_2 w y_1 \dots y_{n-1}))(z_1 y_n) \\ &\stackrel{(2.3)}{\approx} (x(z_2 y_1 \dots y_{n-1} w))(z_1 y_n) \stackrel{(2.7)}{\approx} (x(z_2 y_1 \dots y_n))(z_1 w). \end{aligned}$$

5. The identity (3.12) follows directly by (3.2).

6. The identity (3.13) follows by (2.3), (2.7) and (3.11).  $\square$

The next theorem describes all linear terms in the variety  $\mathcal{V}_{A(P_2+L_2)}$ .

**Theorem 3.4.** *Every term in  $\mathcal{V}_{A(P_2+L_2)}$  may be expressed in one of the following standard forms:*

$$\begin{aligned} \text{(I1)} \quad &xy_1 \dots y_k(t_1 w_1) \dots (t_n w_n), \quad k \geq 0, n \geq 0; \\ \text{(I2)} \quad &(x(y_1 y_2 \dots y_k))(t_1 w_1) \dots (t_n w_n), \quad k \geq 2, n \geq 0. \end{aligned}$$

*Proof.* We show (using  $\Sigma$ ) that the set of terms of the form (I1) and (I2) is closed under multiplication.

a) Let  $\gamma \approx xy_1 \dots y_k(t_1 w_1) \dots (t_n w_n)$  and  $\delta \approx ab_1 \dots b_l(c_1 d_1) \dots (c_m d_m)$ . By (3.3) and (3.4) we have

$$\begin{aligned} \gamma\delta &\approx [xy_1 \dots y_k(t_1 w_1) \dots (t_n w_n)][ab_1 \dots b_l(c_1 d_1) \dots (c_m d_m)] \\ &\stackrel{(3.3)}{\approx} xy_1 \dots y_k(ab_1 \dots b_l(c_1 d_1) \dots (c_m d_m))(t_1 w_1) \dots (t_n w_n) \\ &\stackrel{(3.4)}{\approx} ((xy_1 \dots y_k)(ab_1 \dots b_l))(d_1 c_1) \dots (d_m c_m)(t_1 w_1) \dots (t_n w_n). \end{aligned}$$

If  $l > k \geq 0$ , then

$$\begin{aligned} &((xy_1 \dots y_k)(ab_1 \dots b_l))(d_1 c_1) \dots (d_m c_m)(t_1 w_1) \dots (t_n w_n) \\ &\stackrel{(3.12)}{\approx} (x(ab_1 \dots b_{l-k}))(y_1 b_{l-k+1}) \dots (y_k b_l)(d_1 c_1) \dots (d_m c_m)(t_1 w_1) \dots (t_n w_n). \end{aligned}$$

Hence  $\gamma\delta$  can be written in the form (I2).

For  $k \geq l \geq 0$ ,

$$\begin{aligned} &((xy_1 \dots y_k)(ab_1 \dots b_l))(d_1 c_1) \dots (d_m c_m)(t_1 w_1) \dots (t_n w_n) \\ &\stackrel{(3.12)}{\approx} xy_1 \dots y_{k-l} a(y_{k-l+1} b_1) \dots (y_k b_l)(d_1 c_1) \dots (d_m c_m)(t_1 w_1) \dots (t_n w_n), \end{aligned}$$

and  $\gamma\delta$  has the form (I1).

b) Let  $\gamma \approx xy_1 \dots y_k(t_1 w_1) \dots (t_n w_n)$  and  $\delta \approx (a(b_1 b_2 \dots b_l))(c_1 d_1) \dots (c_m d_m)$  for  $l \geq 2$ . Then

$$\begin{aligned} \gamma\delta &\approx [xy_1 \dots y_k(t_1 w_1) \dots (t_n w_n)][(a(b_1 b_2 \dots b_l))(c_1 d_1) \dots (c_m d_m)] \\ &\stackrel{(3.3)}{\approx} x((a(b_1 b_2 \dots b_l))(c_1 d_1) \dots (c_m d_m))y_1 \dots y_k(t_1 w_1) \dots (t_n w_n) \\ &\stackrel{(3.4),(3.3)}{\approx} x(a(b_1 b_2 \dots b_l))y_1 \dots y_k(d_1 c_1) \dots (d_m c_m)(t_1 w_1) \dots (t_n w_n). \end{aligned}$$

By (3.8) we have that  $x(a(b_1 b_2 \dots b_l)) \approx xab_3 \dots b_l(b_2 b_1)$ . Consequently

$$\begin{aligned} &x(a(b_1 b_2 \dots b_l))y_1 \dots y_k(d_1 c_1) \dots (d_m c_m)(t_1 w_1) \dots (t_n w_n) \\ &\approx xab_3 \dots b_l y_1 \dots y_k(b_2 b_1)(d_1 c_1) \dots (d_m c_m)(t_1 w_1) \dots (t_n w_n), \end{aligned}$$

and  $\gamma\delta$  is reduced to the form (I1).

c) Let  $\gamma \approx (a(b_1 b_2 \dots b_l))(c_1 d_1) \dots (c_m d_m)$  and  $\delta \approx xy_1 \dots y_k(t_1 w_1) \dots (t_n w_n)$ , for  $l \geq 2$ . Then

$$\begin{aligned} \gamma\delta &\approx [(a(b_1 b_2 \dots b_l))(c_1 d_1) \dots (c_m d_m)][xy_1 \dots y_k(t_1 w_1) \dots (t_n w_n)] \\ &\stackrel{(3.3)}{\approx} (a(b_1 b_2 \dots b_l))(xy_1 \dots y_k(t_1 w_1) \dots (t_n w_n))(c_1 d_1) \dots (c_m d_m) \\ &\stackrel{(3.4)}{\approx} (a(b_1 b_2 \dots b_l))(xy_1 \dots y_k)(w_1 t_1) \dots (w_n t_n)(c_1 d_1) \dots (c_m d_m). \end{aligned}$$

If  $k = 0$ , then by (3.10)

$$\gamma\delta \approx (a(b_1 b_2 \dots b_{l-1}))(x b_l)(w_1 t_1) \dots (w_n t_n)(c_1 d_1) \dots (c_m d_m).$$

If  $k \geq 1$ , then by (3.9) we have

$$(a(b_1 b_2 \dots b_l))(xy_1 \dots y_k) \stackrel{(3.9)}{\approx} (a(b_1 b_2 \dots b_l y_2 \dots y_k))(xy_1).$$

It follows that

$$\begin{aligned} &(a(b_1 b_2 \dots b_l))(xy_1 \dots y_k)(w_1 t_1) \dots (w_n t_n)(c_1 d_1) \dots (c_m d_m) \\ &\approx (a(b_1 b_2 \dots b_l y_2 \dots y_k))(xy_1)(w_1 t_1) \dots (w_n t_n)(c_1 d_1) \dots (c_m d_m). \end{aligned}$$

Hence, for any  $k \geq 0$ ,  $\gamma\delta$  may be reduced to the form (I2).

d) Let  $\gamma \approx (x(y_1 y_2 \dots y_k))(t_1 w_1) \dots (t_n w_n)$  and  $\delta \approx (a(b_1 b_2 \dots b_l))(c_1 d_1) \dots (c_m d_m)$ , for  $k, l \geq 2$ . By (3.4) and (E), we have

$$\begin{aligned} \gamma\delta &\approx [(x(y_1 y_2 \dots y_k))(t_1 w_1) \dots (t_n w_n)][(a(b_1 b_2 \dots b_l))(c_1 d_1) \dots (c_m d_m)] \\ &\stackrel{(3.4)}{\approx} (x((y_1 \dots y_k)(w_1 t_1) \dots (w_n t_n)))(a((b_1 b_2 \dots b_l)(d_1 c_1) \dots (d_m c_m))) \\ &\stackrel{(E)}{\approx} xa(((y_1 \dots y_k)(w_1 t_1) \dots (w_n t_n))((b_1 b_2 \dots b_l)(d_1 c_1) \dots (d_m c_m))). \end{aligned}$$

Thus by (a) and (b), also in this case,  $\gamma\delta$  reduces to the required form.  $\square$

**Proposition 3.5.** *The set  $\Sigma$  is a basis for all linear identities true in the variety of entropic graph algebras.*

*Proof.* According to results of [1], each identity which is satisfied in  $\mathcal{V}_{A(P_2+L_2)}$  is regular. Let  $p \approx q$  be linear and regular identity true in  $\mathcal{V}_{A(P_2+L_2)}$ . By Theorem 3.4, the terms  $p$  and  $q$  are in one of the two standard forms.

*Case 1.* Let  $p$  and  $q$  be of type (I1). There are distinct variables

$\{x, y_1, \dots, y_k, y_{k+1}, \dots, y_{k+n}, z_1, \dots, z_n\} = \{a, b_1, \dots, b_l, b_{l+1}, \dots, b_{l+m}, c_1, \dots, c_m\}$ , such that

$$p \approx xy_1 \dots y_k (y_{k+1} z_1) \dots (y_{k+n} z_n)$$

and

$$q \approx ab_1 \dots b_l (b_{l+1} c_1) \dots (b_{l+m} c_m).$$

By (3.6),

$$p \approx x(y_1 x) \dots (y_k x) (y_{k+1} z_1) \dots (y_{k+n} z_n)$$

and

$$q \approx a(b_1 a) \dots (b_l a) (b_{l+1} c_1) \dots (b_{l+m} c_m).$$

By Theorem 2.1, the identity  $p \approx q$  is satisfied in  $\mathcal{V}_{A(P_2+L_2)}$  if and only if  $x = a$ ,  $\{x, z_1, \dots, z_n\} = \{a, c_1, \dots, c_m\}$  and  $\{y_1, \dots, y_{k+n}\} = \{b_1, \dots, b_{l+m}\}$ . This implies that  $n = m$ ,  $k = l$  and there are permutations  $\pi$  on  $\{1, 2, \dots, k+n\}$  and  $\delta$  on  $\{1, 2, \dots, n\}$  such that

$$q \approx xy_{\pi(1)} \dots y_{\pi(k)} (y_{\pi(k+1)} z_{\delta(1)}) \dots (y_{\pi(k+n)} z_{\delta(n)}).$$

Consequently, by identity (3.5), the identity  $p \approx q$  follows from  $\Sigma$ .

*Case 2.* Let  $p$  and  $q$  be of type (I2). There are distinct variables

$\{x, z_1, \dots, z_{n+1}, y_1, \dots, y_{k+n}\} = \{a, b_1, \dots, b_{m+1}, c_1, \dots, c_{l+m}\}$ , such that

$$p \approx (x(z_1 y_1 \dots y_k))(z_2 y_{k+1}) \dots (z_{n+1} y_{k+n}),$$

$$q \approx (a(b_1 c_1 \dots c_l))(b_2 c_{l+1}) \dots (b_{m+1} c_{l+m}),$$

and  $k, l \geq 2$ .

By the identity (3.7),

$$p \approx x(z_1 x)(z_1 y_1) \dots (z_1 y_k)(z_2 y_{k+1}) \dots (z_{n+1} y_{k+n})$$

and

$$q \approx a(b_1 a)(b_1 c_1) \dots (b_l c_l)(b_{l+1} c_{l+1}) \dots (b_{m+1} c_{l+m}).$$

By Theorem 2.1, the identity  $p \approx q$  is satisfied in  $\mathcal{V}_{A(P_2+L_2)}$  if and only if  $x = a$ ,  $\{x, y_1, \dots, y_{k+n}\} = \{a, c_1, \dots, c_{l+m}\}$  and  $\{z_1, \dots, z_{n+1}\} = \{b_1, \dots, b_{m+1}\}$ . This



implies that  $n = m$ ,  $k = l$  and there are permutations  $\delta$  on  $\{1, 2, \dots, k + n\}$  and  $\pi$  on  $\{1, 2, \dots, n + 1\}$  such that

$$q \approx (x(z_{\pi(1)}y_{\delta(1)} \cdots y_{\delta(k)}))(z_{\pi(2)}y_{\delta(k+1)} \cdots (z_{\pi(n+1)}y_{\delta(k+n)}).$$

Hence, by identity (3.13), the identity  $p \approx q$  follows from  $\Sigma$ .

*Case 3.* Let  $p$  be of type (I1) and  $q$  be of type (I2). As before,

$$p \approx xy_1 \cdots y_k(y_{k+1}z_1) \cdots (y_{k+n}z_n)$$

and

$$q \approx (a(b_1c_1 \cdots c_l))(b_2c_{l+1}) \cdots (b_{m+1}c_{l+m}),$$

where  $\{x, z_1, \dots, z_n, y_1, \dots, y_{k+n}\} = \{a, b_1, \dots, b_{m+1}, c_1, \dots, c_{l+m}\}$  and  $l \geq 2$ . Similarly as in Case 1 and Case 2, the identities (3.6), (3.7) and Theorem 2.1 imply that, the identity  $p \approx q$  is satisfied in  $\mathcal{V}_{A(P_2+L_2)}$  if and only if  $x = a$ ,  $\{x, z_1, \dots, z_n\} = \{a, c_1, \dots, c_{l+m}\}$  and  $\{y_1, \dots, y_{k+n}\} = \{b_1, \dots, b_{m+1}\}$ . It follows that  $n = m + l$  and  $m + 1 = k + n$ . Hence  $k + l = 1$ . But we assumed that  $l \geq 2$ , so this gives a contradiction.

This shows that any linear identity holding in  $\mathcal{V}_{A(P_2+L_2)}$  is a consequence of  $\Sigma$ .  $\square$

#### 4. THE VARIETY OF POWER ENTROPIC GRAPH ALGEBRAS

Using the result of the previous section we will describe the lattice of all subvarieties of power entropic graph algebras.

Let  $\mathcal{V}$  be a variety. We will denote by  $\mathbf{Cm}\mathcal{V}$  the variety generated by power algebras of algebras in  $\mathcal{V}$ , i.e.,

$$\mathbf{Cm}\mathcal{V} := \text{HSP}(\{\mathbf{CmA} \mid \mathbf{A} \in \mathcal{V}\}).$$

Evidently,  $\mathcal{V} \subseteq \mathbf{Cm}\mathcal{V}$ , because every algebra  $\mathbf{A}$  in  $\mathcal{V}$  embeds into  $\mathbf{CmA}$  by  $x \mapsto \{x\}$ . G. Grätzer and H. Lakser proved the following theorem.

**Theorem 4.1.** [5] *Let  $\mathcal{V}$  be a variety. The variety  $\mathbf{Cm}\mathcal{V}$  satisfies precisely those identities resulting through identification of variables from the linear identities true in  $\mathcal{V}$ .*

For example, an idempotent law is satisfied in the variety  $\mathbf{Cm}\mathcal{V}$  if and only if it is a consequence of linear identities true in  $\mathcal{V}$ .

As an immediate corollary of the Theorem 4.1 we have the following result.

**Corollary 4.2.** [5] *For a variety  $\mathcal{V}$ ,  $\mathbf{Cm}\mathcal{V} = \mathcal{V}$  if and only if  $\mathcal{V}$  is defined by a set of linear identities.*

As it was shown in Section 3, any linear identity true in  $\mathcal{V}_{A(P_2+L_2)}$  is derivable from the set  $\Sigma$ . Hence, by Theorem 4.1, we obtain the following result.

**Proposition 4.3.**  $\mathbf{Cm}\mathcal{V}_{A(P_2+L_2)} = \text{Mod}(\Sigma)$ .

Similarly as in the case of the variety  $\mathcal{V}_{A(P_2+L_2)}$  we can show that varieties  $\mathbf{Cm}\mathcal{V}_{A(P_3)}$ ,  $\mathbf{Cm}\mathcal{U}_2$  and  $\mathbf{Cm}\mathcal{V}_{A(P_3+L_1)}$  are also defined by the set  $\Sigma$ .

Now we will give a description of varieties generated by power algebras from remaining subvarieties of  $\mathcal{V}_{A(P_2+L_2)}$ . The varieties  $\mathcal{T}$ ,  $\mathcal{LZ}$  and  $\mathcal{V}_{A(P_1)}$  are defined only by linear identities. Hence by Corollary 4.2,  $\mathbf{Cm}\mathcal{T} = \mathcal{T}$ ,  $\mathbf{Cm}\mathcal{LZ} = \mathcal{LZ}$  and  $\mathbf{Cm}\mathcal{V}_{A(P_1)} = \mathcal{V}_{A(P_1)}$ .

The varieties  $\mathcal{SL}$  and  $\mathcal{V}_{A(P_1+L_1)}$  satisfy associativity and commutativity. Hence,  $\mathbf{Cm}\mathcal{SL} = \mathbf{Cm}\mathcal{V}_{A(P_1+L_1)}$  is the variety  $\mathcal{CS}$  of commutative semigroups.

It is easy to see that by left-normal law (2.3) and associativity, a linear regular identity  $p \approx q$  is true in the variety  $\mathcal{LN}$  or  $\mathcal{V}_{A(P_1+L_2)}$  if and only if terms  $p$  and  $q$  have the same leftmost variable. This implies that  $\mathbf{Cm}\mathcal{LN} = \mathbf{Cm}\mathcal{V}_{A(P_1+L_2)}$  is the variety  $\mathcal{LS}$  of left-normal semigroups.

Moreover, the variety  $\mathcal{U}_1$  satisfies the associative law. This implies that each linear identity in  $\mathcal{U}_1$  is a consequence of the identity  $xy \approx xz$  and associativity. Hence  $\mathbf{Cm}\mathcal{U}_1$  coincides with the variety  $\mathcal{US}$  defined by  $xy \approx xz$  and associativity.

Now let us consider the following linear identity:

$$(4.1) \quad x(yz) \approx z(yx).$$

It is not difficult to see that the identity (3.1) is a consequence of entropy, left-normality and the identity (4.1):

$$x(y(zt)) \stackrel{(4.1)}{\approx} zt(yx) \stackrel{(2.3)}{\approx} z(yx)t \stackrel{(4.1)}{\approx} x(yz)t \stackrel{(2.3)}{\approx} xt(yz) \stackrel{(E)}{\approx} xy(tz).$$

Let  $\Gamma$  be the following set of identities:

$$\Gamma : \begin{cases} (E) & xy(zt) \approx xz(yt), \\ (2.3) & xyz \approx xzy, \\ (4.1) & x(yz) \approx z(yx). \end{cases}$$

**Lemma 4.4.** *For  $n \geq 1$ , the identity*

$$(4.2) \quad x_1y_1(y_2x_2) \dots (y_nx_n) \approx x_{\pi(1)}y_{\delta(1)}(y_{\delta(2)}x_{\pi(2)}) \dots (y_{\delta(n)}x_{\pi(n)})$$

*follows from the set  $\Gamma$  for any permutations  $\pi$  and  $\delta$  of the set  $\{1, 2, \dots, n\}$ .*

*Proof.* This follows directly by (3.5) and (4.1).  $\square$

**Theorem 4.5.** *Any linear identity holding in the variety  $\mathcal{V}_{A(P_2+L_1)}$  is a consequence of the set  $\Gamma$ .*

*Proof.* Using the same methods as in proofs of Theorem 3.4 and Proposition 3.5 we obtain that each linear regular identity true in  $\mathcal{V}_{A(P_2+L_1)}$  is one of the two following types:

$$\begin{aligned} & x_1y_1 \dots y_k(y_{k+1}x_2) \dots (y_{n+k}x_{n+1}) \\ & \approx x_{\delta(1)}y_{\pi(1)} \dots y_{\pi(k)}(y_{\pi(k+1)}x_{\delta(2)}) \dots (y_{\pi(n+k)}x_{\delta(n+1)}), \end{aligned}$$

for any permutation  $\pi$  on the set  $\{1, \dots, n+k\}$  and any permutation  $\delta$  on the set  $\{1, \dots, n+1\}$ , or

$$\begin{aligned} & (x_1(y_1x_2 \dots x_k))(y_2x_{k+1}) \dots (y_{n+1}x_{k+n}) \\ & \approx (x_{\delta(1)}(y_{\pi(1)}x_{\delta(2)} \dots x_{\delta(k)}))(y_{\pi(2)}x_{\delta(k+1)}) \dots (y_{\pi(n+1)}x_{\delta(k+n)}), \end{aligned}$$

for any permutation  $\pi$  on the set  $\{1, \dots, n+1\}$  and any permutation  $\delta$  on the set  $\{1, \dots, n+k\}$ .

By identities (3.5), (3.13), (4.1) and (4.2), both follow from  $\Gamma$ .  $\square$

Similarly, we can show that any linear identity true in  $\mathcal{V}_{A(P_2)}$  is also a consequence of  $\Gamma$ . All these observations prove the following proposition.

**Proposition 4.6.** *There are eight varieties of power entropic graph algebras:*

- $\mathbf{Cm}\mathcal{V}_{A(P_2+L_2)} = \mathbf{Cm}\mathcal{V}_{A(P_3)} = \mathbf{Cm}\mathcal{U}_2 = \mathbf{Cm}\mathcal{V}_{A(P_3+L_1)} = \text{Mod}(\Sigma)$
- $\mathbf{Cm}\mathcal{V}_{A(P_2+L_1)} = \mathbf{Cm}\mathcal{V}_{A(P_2)} = \text{Mod}(\Gamma)$
- $\mathbf{Cm}\mathcal{L}\mathcal{N} = \mathbf{Cm}\mathcal{V}_{A(P_1+L_2)} = \mathcal{L}\mathcal{S} = \text{Mod}(xyz \approx xzy, xyz \approx x(yz))$
- $\mathbf{Cm}\mathcal{S}\mathcal{L} = \mathbf{Cm}\mathcal{V}_{A(P_1+L_1)} = \mathcal{C}\mathcal{S} = \text{Mod}(xy \approx yx, xyz \approx x(yz))$
- $\mathbf{Cm}\mathcal{U}_1 = \mathcal{U}\mathcal{S} = \text{Mod}(xy \approx xz, xyz \approx x(yz))$
- $\mathbf{Cm}\mathcal{V}_{A(P_1)} = \mathcal{V}_{A(P_1)} = \text{Mod}(xy \approx uz)$
- $\mathbf{Cm}\mathcal{L}\mathcal{Z} = \mathcal{L}\mathcal{Z} = \text{Mod}(xy \approx x)$
- $\mathbf{Cm}\mathcal{T} = \mathcal{T} = \text{Mod}(x \approx y)$

The lattice of all subvarieties of  $\mathbf{Cm}\mathcal{V}_{A(P_2+L_2)}$  is given in the Figure 2:




FIGURE 2

REFERENCES

- [1] K.A. Baker, G.F. McNulty, H.Werner, *The finitely based varieties of graph algebras*, Acta Scient. Math. 51 (1987), 3–15.
- [2] I. Bošnjak, R. Madarász, *On power structures*, Algebra and Discr. Math. 2 (2003), 14–35.
- [3] C. Brink, *Power structures*, Algebra Univers. 30 (1993), 177–216.
- [4] B.A. Davey, P.H. Idziak, W.A. Lampe, G.F. McNulty, *Dualizability and graph algebras*, Discrete Math. 214 (2000), 145–172.
- [5] G. Grätzer, H. Lakser, *Identities for globals (complex algebras) of algebras*, Colloq. Math. 56 (1988), 19–29.
- [6] G. Grätzer, S. Whitney, *Infinitary varieties of structures closed under the formation of complex structures*, Colloq. Math. 48 (1984), 485–488.
- [7] G.F. McNulty, C. Shallon, *Inherently nonfinitely based finite algebras*, R.Freese, O.Garcia (Eds.), Universal Algebra and Lattice Theory, Proceedings, Puebla 1982, Lecture Notes in Mathematics, Springer, Berlin, 1004 (1983), 205–231.
- [8] A. Shafaat, *On varieties closed under the construction of power algebras*, Bull. Austral. Math. Soc. 11 (1974), 213–218.
- [9] C.R. Shallon, *Nonfinitely based binary algebras derived from lattices*, Ph.D. Thesis, University of California at Los Angeles, 1979.

FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, WARSAW UNIVERSITY OF TECHNOLOGY, PLAC POLITECHNIKI 1, 00-661 WARSAW, POLAND

E-mail address: apili@mini.pw.edu.pl

URL: <http://www.mini.pw.edu.pl/~apili>